arXiv:1604.06930v1 [math.NT] 23 Apr 2016

THREE STAGE TOWERS OF 5-CLASS FIELDS

DANIEL C. MAYER

ABSTRACT. With $K = \mathbb{Q}\left(\sqrt{3\,812\,377}\right)$ we give the first example of an algebraic number field possessing a 5-class tower of exact length $\ell_5 K = 3$. The rigorous proof is conducted by means of the p-group generation algorithm, showing the existence of a unique finite metabelian 5-group $\mathfrak{G}$ with abelianization $[5,5]$ having the kernels $(M_1, \mathfrak{G}^5)$ and targets $([25,5,5,5],[5,5]^5)$ of Artin transfers $T_i : \mathfrak{G} \to M_i/M_i'$ to its six maximal subgroups M_i, prescribed by arithmetical invariants of K. Thus, $\mathfrak{G}$ must be the second 5-class group $\mathrm{G}_5^2 K$ of the real quadratic field K but cannot be its 5-class tower group $\mathrm{G}_5^\infty K$, since the relation rank $d_2\mathfrak{G} = 4$ is too big. We provide evidence of exactly five non-isomorphic extensions G of $\mathfrak{G}$ having the required relation rank $d_2 G = 3$ and derived length $\mathrm{dl}(G) = 3$ whose metabelianization G/G'' is isomorphic to $\mathfrak{G}$. Consequently, $\mathrm{G}_5^\infty K$ must be one of the five non-metabelian groups G.

1. INTRODUCTION

Given a prime number $p \geq 2$, a positive integer $n \geq 1$, and an algebraic number field K, the nth *Hilbert p-class field* $\mathrm{F}_p^n K$ of K is the maximal unramified Galois extension of K whose automorphism group $\mathrm{G}_p^n K := \mathrm{Gal}\left(\mathrm{F}_p^n K | K\right)$ is a finite p-group of derived length $\mathrm{dl}(\mathrm{G}_p^n K) \leq n$. In particular, the first Hilbert p-class field $\mathrm{F}_p^1 K$ of K is the maximal abelian unramified p-extension of K. Its Galois group $\mathrm{G}_p^1 K$ is isomorphic to the p-class group $\mathrm{Cl}_p K := \mathrm{Syl}_p \mathrm{Cl}(K)$ of K, according to Artin's reciprocity law of class field theory [1].

The projective limit $\varprojlim_{n \geq 1} \mathrm{G}_p^n K$ of all nth *p-class groups* $\mathrm{G}_p^n K$ of K is isomorphic to the possibly infinite topological Galois group $G := \mathrm{G}_p^\infty K$ of the maximal unramified pro-p extension $\mathrm{F}_p^\infty K := \bigcup_{n \geq 1} \mathrm{F}_p^n K$, that is the *Hilbert p-class tower*, of K. Conversely, the nth p-class group can be obtained as the nth derived quotient $G/G^{(n)} \simeq \mathrm{G}_p^n K$ of the *p-class tower group* G of K.

The *length* of the p-class tower of K is defined as the derived length of G,

$$\ell_p K := \mathrm{dl}\left(\mathrm{G}_p^\infty K\right). \tag{1.1}$$

For each $n \geq 2$, inclusively $n = \infty$, the commutator subgroup of $\mathrm{G}_p^n K$ is isomorphic to $\mathrm{Gal}\left(\mathrm{F}_p^n K | \mathrm{F}_p^1 K\right)$, and the abelianization of $\mathrm{G}_p^n K$ is isomorphic to the p-class group $\mathrm{Cl}_p K$ of K,

$$\mathrm{G}_p^n K / \left(\mathrm{G}_p^n K\right)' \simeq \mathrm{G}_p^1 K \simeq \mathrm{Cl}_p K, \tag{1.2}$$

in particular, the generator rank $d_1\left(\mathrm{G}_p^n K\right) := \dim_{\mathbb{F}_p} \mathrm{H}^1(\mathrm{G}_p^n K, \mathbb{F}_p)$ of $\mathrm{G}_p^n K$ coincides with the p-class rank $r_p K := \dim_{\mathbb{F}_p}\left(\mathrm{Cl}_p K \bigotimes_{\mathbb{Z}_p} \mathbb{F}_p\right)$ of K, according to Burnside's basis theorem.

After the preceding clarification of the technical framework for this article, we devote a minimum of space to an outline of the historical evolution of finding p-class towers with increasing finite length. Thereby, we use the notation of the SmallGroups database [4], [5], and partially give more details than the original papers of the cited authors,

We have $\ell_p K = 0$ if and only if the class number $\#\mathrm{Cl}(K)$ of the field K is not divisible by p.

Date: April 23, 2016.
2000 *Mathematics Subject Classification.* Primary 11R37, 11R29, 11R11, 11R20; Secondary 20D15, 20–04.
Key words and phrases. 5-class towers, 5-class groups, 5-capitulation, real quadratic fields, dihedral fields of degree 10; finite 5-groups with two generators, descendant tree, p-group generation algorithm, nuclear rank, bifurcation, p-multiplicator rank, relation rank, generator rank, Shafarevich cover, Artin transfers, partial order of Artin patterns.
Research supported by the Austrian Science Fund (FWF): P 26008-N25.

If the p-class group $\mathrm{Cl}_p K > 1$ is cyclic, then we have a single stage tower with $\ell_p K = 1$, by Formula (1.2). If $K = \mathbb{Q}\left(\sqrt{d}\right)$ is a quadratic field and $p \geq 3$ is odd, the converse is also true [23, Thm. 4.1, (1), p. 486]. However, for the discriminant $d = -84 = -2^2 \cdot 3 \cdot 7$, we have $\mathrm{Cl}_2 K \simeq [2,2]$ but nevertheless only $\ell_2 K = 1$ [20, (ii), p. 277], [23, § 9, pp. 501–503].

In 1930, Hasse [17, § 27, pp. 173–174] gave the first example of $\ell_2 K \geq 2$ for a field of type $[4,2]$, due to private communications by Furtwängler and Artin. It has discriminant $d = -260 = -2^2 \cdot 5 \cdot 13$, $\mathrm{G}_2^2 K \simeq \langle 2^4, 6 \rangle$, and in fact $\ell_2 K = 2$. A field of type $[2,2]$ with $\ell_2 K = 2$ is given by $d = -120 = -2^3 \cdot 3 \cdot 5$ with $\left(\frac{5}{3}\right) = -1$, and has $\mathrm{G}_2^2 K \simeq \langle 2^3, 4 \rangle$ the quaternion group [20, (vi.a,b), p. 278]. However, a field of type $[2,2]$ can never have a 2-class tower of length bigger than two.

Consequently, the first field with $\ell_2 K = 3$ was of type $[4,2]$. It was discovered in 2003 by Bush [10, Prop. 2, p. 321] and has discriminant $d = -1\,780 = -2^2 \cdot 5 \cdot 89$. Its 2-tower group is one of two candidates $\langle 2^8, 426|427 \rangle$ with class 5 and coclass 3. In 2006, Nover provided evidence of $\ell_2 K = 3$ for a field of type $[2,2,2]$ with $d = -3\,135 = -3 \cdot 5 \cdot 11 \cdot 19$ and four candidates for $\mathrm{G}_2^3 K$ of order 2^{13}, class 7 and coclass 6 [9, § 3.4, pp. 7–8], [34, Prop. 5, p. 239].

Now we turn to odd primes $p \geq 3$. In 1934, Scholz and Taussky [37, § 3, pp. 39–41] manually calculated the first occurrences of $\ell_3 K = 2$ for $d = -4\,027$ of type $[3,3]$, resp. $d = -3\,299$ of type $[9,3]$. The corresponding 3-class tower groups are the Schur σ-groups $\langle 3^5, 5 \rangle$, resp. one of the two candidates $\langle 3^6, 14|15 \rangle$. However, they also claimed a two stage tower for $d = -9\,748 = 2^2 \cdot 2\,437$ [37, § 3, p. 41], where neither of the two candidates $\langle 3^7, 302|306 \rangle$ for the second 3-class group is a Schur σ-group [21]. This claim was corrected by Bush and ourselves [11, Cor. 4.1.1, p. 775] in 2015. We proved the first sufficient criterion for complex quadratic fields K with $\ell_3 K = 3$, and provided two candidates $\mathrm{G}_3^3 K \simeq \langle 3^6, 54 \rangle - \#2; 2|6$ for $d = -9\,748$, in the notation of the ANUPQ package [14]. These were the first non-metabelian Schur σ-groups. They have order 3^8, class 5, coclass 3, derived length 3, and are discussed in more detail in [26, (70), p. 190], [26, Fig. 10, p. 191], and [28, §§ 6–7, p. 751–756].

The first occurrences of $\ell_5 K = 2$, resp. $\ell_7 K = 2$, were discovered in 2013 by ourselves for $d = -11\,199$ [25, Tbl. 3.13, p. 450], resp. $d = -63\,499$ [25, Tbl. 3.14, p. 450]. The corresponding groups $\mathrm{G}_5^2 K \simeq \langle 5^5, 12 \rangle$, resp. $\mathrm{G}_7^2 K \simeq \langle 7^5, 14|16 \rangle$ with two candidates, are metabelian Schur σ-groups. Our attempt to prove $\ell_5 K = 3$ for $d = -62\,632$ with two candidates $\mathrm{G}_5^2 K \simeq \langle 5^6, 564 \rangle - \#1; 2|3$, resp. $d = -67\,063$ with six candidates $\mathrm{G}_5^2 K \simeq \langle 5^6, 564 \rangle - \#1; 8|9|11|12|14|15$, was not successful, since we could not find suitable Schur σ-groups $\mathrm{G}_5^3 K$.

Little progress was achieved for real quadratic fields until we developed new techniques for determining their p-class tower in [27]. We found $\ell_3 K = 2$ for $d = 422\,573$ with $\mathrm{G}_3^2 K \simeq \langle 3^5, 5 \rangle$ [23, Tbl. 4, p. 498], and $\ell_5 K = 2$ for $d = 4\,954\,652$ with $\mathrm{G}_5^2 K \simeq \langle 5^5, 12 \rangle$. With extensive computational effort we succeeded in proving $\ell_3 K = 3$ for $d = 957\,013$, which has $\mathrm{G}_3^3 K \simeq \langle 3^7, 273 \rangle$ with class 5, coclass 2, and relation rank 3 [27, Exm. 6.3, pp. 306–307]. However, the attempt to prove that the quintic analogue $d = 12\,562\,849$ has $\ell_5 K = 3$ with either of the two candidates $\langle 5^6, 680 \rangle - \#1; 1|2$ for $\mathrm{G}_5^3 K$, having order 5^7, class 5 and coclass 2, failed, due to a complete exhaustion of RAM during the construction of two successive unramified cyclic quintic extensions of the real quadratic field K, that is an extension of absolute degree 50.

Exploring p-groups of coclass 1 instead of coclass 2, which would be useless for finding 3-class towers of three stages, turned out to be the crucial idea how to circumvent such hopeless number theoretic computations of high complexity and to conduct a purely group theoretic proof of the principal results of this article, which are the following statements, in a succinct coarse form.

Theorem 1.1. *Let K be a real quadratic field with 5-class group $\mathrm{Cl}_5 K$ of type $[5,5]$ and denote by $L_1, \ldots, L_6$ its six unramified cyclic quintic extensions. If K possesses the 5-capitulation type*

$$\varkappa(K) \sim (1,0,0,0,0,0), \text{ with fixed point } 1, \tag{1.3}$$

in the six extensions L_i, and if the 5-class groups $\mathrm{Cl}_5 L_i$ are given by

$$\tau(K) \sim ([25,5,5,5],[5,5],[5,5],[5,5],[5,5],[5,5]), \tag{1.4}$$

then the 5-class tower $K < \mathrm{F}_5^1K < \mathrm{F}_5^2K < \mathrm{F}_5^3K = \mathrm{F}_5^\infty K$ *of* K *has exact length* $\ell_5K = 3$.

Corollary 1.1. *A real quadratic field* K *which satisfies the assumptions in Theorem 1.1, in particular the Formulas* (1.3) *and* (1.4)*, has the unique second* 5*-class group*

$$\mathrm{G}_5^2K \simeq \langle 15625, 635\rangle \tag{1.5}$$

with order 5^6*, class* 5*, coclass* 1*, derived length* 2*, and relation rank* 4*, and one of the following five candidates for the* 5*-class tower group*

$$\mathrm{G}_5^\infty K = \mathrm{G}_5^3K \simeq \langle 78125, n\rangle, \quad \text{where } n \in \{361, 373, 374, 385, 386\}, \tag{1.6}$$

with order 5^7*, class* 5*, coclass* 2*, derived length* 3*, and relation rank* 3.

Example 1.1. The minimal fundamental discriminant d of a real quadratic field $K = \mathbb{Q}(\sqrt{d})$ satisfying the conditions (1.3) and (1.4) is given by

$$d = 3\,812\,377 = 991 \cdot 3\,847. \tag{1.7}$$

The next occurrence is $d = 19\,621\,905 = 3 \cdot 5 \cdot 307 \cdot 4\,261$, which is currently the biggest known example, whereas $d = 21\,281\,673 = 3 \cdot 7 \cdot 53 \cdot 19\,121$ satisfies (1.4) but has a different $\varkappa(K) \sim (2, 0, 0, 0, 0, 0)$, without fixed point.

The organization of this paper is as follows. In § 2, we show how the arithmetical invariants $\varkappa(K)$ and $\tau(K)$ of an algebraic number field K (e.g., the data in Formula (1.3) and (1.4)) can be translated into group theoretic information $\varkappa(\mathfrak{G})$ and $\tau(\mathfrak{G})$ on the second p-class group $\mathfrak{G} = \mathrm{G}_p^2K$ of K, following the ideas indicated in Artin's paper [2], which led to Furtwängler's proof [13] of the famous Principal Ideal Theorem and were expanded in detail by Hasse [17].

§ 3 is devoted to the study of the interplay between the Artin pattern $\mathrm{AP}(G) = (\tau(G), \varkappa(G))$ of finite p-groups G on a descendant tree $\mathcal{T}(R)$ with root R and the partial order $G > \pi G$ induced by the (child, parent)-pairs $(G, \pi G)$ of the tree, which was discovered recently in [30]. These monotony properties of $\tau(G)$ and $\varkappa(G)$ are employed in the p-group generation algorithm by Newman [33] and O'Brien [35] for identifying the unique metabelianization $\mathfrak{G} = \mathrm{G}_5^2K = G/G'' \simeq \langle 15625, 635\rangle$ of the 5-class tower group $G = \mathrm{G}_5^\infty K$ of K by the strategy of pattern recognition via Artin transfers in § 4. An unexpected bifurcation in the tree $\mathcal{T}(\langle 25, 2\rangle)$ enabled the discovery of five non-isomorphic possibilities $G \simeq \langle 78125, n\rangle$ with $n \in \{361, 373, 374, 385, 386\}$ and derived length $\mathrm{dl}(G) = 3$ for G itself.

The termination condition for the algorithm is expressed by the growth of the $\tau(G)$-component beyond a break-off bound. At this stage it is still open whether the length of the 5-class tower of K is $\ell_5K = 3$ or $\ell_5K = 2$, since only group theoretic information was used up to this point.

Therefore, arithmetical constraints for G in form of the relation rank, which has been determined by Shafarevich [36], come into the play in §§ 5 and 6, and eliminate the possibility of a metabelian tower with length $\ell_5K = 2$ for a real quadratic field K.

2. Pattern recognition via Artin transfers

2.1. **Translation from number theory to group theory.** We start with an arithmetical situation where Artin patterns play an important role. Let $p \geq 2$ be a prime number. Suppose we are given an algebraic number field $K|\mathbb{Q}$ with non-trivial p-class group $\mathrm{Cl}_pK > 1$. Then the intermediate fields L between K and its Hilbert p-class field F_p^1K are exactly the (finitely many) abelian unramified extensions of K with degree a power of p. For each of them, we denote by $j_{L|K} : \mathrm{Cl}_pK \to \mathrm{Cl}_pL$ the *class extension* homomorphism.

Definition 2.1. By the *restricted Artin pattern* of K we understand the pair consisting of the family $\tau(K)$ of the p-class groups of all extensions $K < L < \mathrm{F}_p^1K$ as its first component and the *p-capitulation type* $\varkappa(K)$ as its second component,

$$\mathrm{AP}(K) := (\tau(K), \varkappa(K)), \quad \tau(K) := (\mathrm{Cl}_p L)_{K<L<\mathrm{F}_p^1 K}, \quad \varkappa(K) := \left(\ker j_{L|K}\right)_{K<L<\mathrm{F}_p^1 K}. \tag{2.1}$$

Remark 2.1. It is convenient to replace the group objects in the unordered family $\tau(K)$, resp. $\varkappa(K)$, by ordered abelian type invariants, resp. ordered numerical identifiers, accepting the drawback of dependence on the selected ordering (indicated by using the symbol $\sim$ instead of $=$).

In the case $p = 5$ and $C := \mathrm{Cl}_5 K \simeq [5,5]$, which is relevant for this paper, let $L_1, \ldots, L_6$ be the six intermediate fields between K and $\mathrm{F}_5^1 K$ and denote by $N_i := \mathrm{Norm}_{L_i|K}(\mathrm{Cl}_5 L_i)$ their norm class groups, which are exactly the six cyclic subgroups of index 5 in C. In this sense, the notation of the 5-capitulation type in Formula (1.3) (where 0 means a total capitulation) is an abbreviation for

$$\varkappa(K) \sim (N_1, C, C, C, C, C). \tag{2.2}$$

Now we turn to the group theoretic aspect of Artin patterns. Let G be a finite p-group or an infinite topological pro-p group with finite abelianization G/G'. For each of the (finitely many) intermediate groups M between the commutator subgroup G' and the group G, we denote by $T_{G,M} : G \to M/M'$ the *Artin transfer* homomorphism from G to the abelianization M/M' of M [26, Dfn. 3.1, p. 5].

Definition 2.2. By the *restricted Artin pattern* of G we understand the pair consisting of the *transfer target type* (TTT) $\tau(G)$ as its first component and the *transfer kernel type* (TKT) $\varkappa(G)$ as its second component,

$$\mathrm{AP}(G) := (\tau(G), \varkappa(G)), \quad \tau(G) := (M/M')_{G'<M<G}, \quad \varkappa(G) := (\ker T_{G,M})_{G'<M<G}. \tag{2.3}$$

Remark 2.2. Again, we usually replace the group objects in the unordered family $\tau(G)$, resp. $\varkappa(G)$, by ordered abelian type invariants, resp. ordered numerical identifiers.

In the case $p = 5$ and $G/G' \simeq [5,5]$, which will be of concern in this paper, we denote by $M_1, \ldots, M_6$ the six intermediate groups between G' and G and we write the transfer kernel type corresponding to Formula (1.3) as

$$\varkappa(G) \sim (M_1, G, G, G, G, G) \quad \text{or briefly } \varkappa(G) \sim (1,0,0,0,0,0). \tag{2.4}$$

Finally, we combine the number theoretic and group theoretic view of Artin patterns.

Theorem 2.1. *Let p be a prime number. Assume that K is a number field, and let $\mathfrak{G} := \mathrm{G}_p^2 K$ be the second p-class group of K. Then $\mathfrak{G}$ and K share a common restricted Artin pattern,*

$$\mathrm{AP}(\mathfrak{G}) = \mathrm{AP}(K), \quad \textit{that is} \quad \tau(\mathfrak{G}) = \tau(K) \textit{ and } \varkappa(\mathfrak{G}) = \varkappa(K). \tag{2.5}$$

Proof. A sketch of the proof is indicated in [32] and [24, § 2.3, pp. 476–478], but the precise proof has been given by Hasse in [17, § 27, pp. 164–175]. We have elaborated Hasse's proof on Wikipedia:
`https://en.wikipedia.org/wiki/Principalization_(algebra)` □

2.2. **Metabelian p-groups of maximal class.** Generally, let G be a finite p-group, for an arbitrary prime number $p \geq 2$. Assume that G is of order $|G| = p^n$, where n is the *logarithmic order*, of *nilpotency class* $c = \mathrm{cl}(G) = m - 1$, where m is the *index of nilpotency*, and of *coclass* $r = \mathrm{cc}(G) = n - c = n - m + 1$. Denote the descending *lower central series* of G by

$$G = \gamma_1 G > \gamma_2 G > \gamma_3 G > \ldots > \gamma_{m-1} G > \gamma_m G = 1, \tag{2.6}$$

where $\gamma_j G = [\gamma_{j-1} G, G]$, for $j \geq 2$, and the ascending *upper central series* of G by

$$1 = \zeta_0 G < \zeta_1 G < \zeta_2 G < \ldots < \zeta_{c-1} G < \zeta_c G = G, \tag{2.7}$$

where $\zeta_j G/\zeta_{j-1}G = \mathrm{Centre}(G/\zeta_{j-1}G)$, for $j \geq 1$.

Now let G be a metabelian p-group of coclass $r = \mathrm{cc}(G) = 1$, whence $n = m$. Then the abelianization G/G' is of type $[p,p]$ and $G = \langle x, y\rangle$ can be generated by two elements x, y. Suppose that $m \geq 3$, i.e., G is non-abelian. Denote the *two-step centralizer*, that is the centralizer of the two-step factor group $\gamma_2 G/\gamma_4 G$ of the lower central series, by

$$\begin{aligned} \chi_2 G &:= \{g \in G \mid [g,u] \in \gamma_4 G \quad \text{for all } u \in \gamma_2 G\}, \text{ that is,} \\ \chi_2 G/\gamma_4 G &= \mathrm{Centralizer}_{G/\gamma_4 G}(\gamma_2 G/\gamma_4 G). \end{aligned} \tag{2.8}$$

$\chi_2 G$ is the biggest subgroup of G such that $[\chi_2 G, \gamma_2 G] \leq \gamma_4 G$. It is characteristic, contains the commutator subgroup $\gamma_2 G$, and coincides with G if and only if $m = 3$, whence we have $\gamma_2 G < \chi_2 G < G \Longleftrightarrow m \geq 4$. Select *normalized generators* of $G = \langle x, y\rangle$ such that

$$x \in G \setminus \chi_2 G, \text{ if } m \geq 4, \quad y \in \chi_2 G \setminus \gamma_2 G. \tag{2.9}$$

Starting with the main commutator $s_2 := [y, x] \in \gamma_2 G$ define the *higher iterated commutators* recursively by $s_j := [s_{j-1}, x] \in \gamma_j G$, for $j \geq 3$.

The general theory of p-groups of maximal class can be found in Huppert [19, § 14, p. 361] and Berkovich [3, § 9, p. 114].

In the isomorphism class of a metabelian p-group G of maximal class, and of order $|G| = p^m$, there exists a representative $G_a^m(z,w)$ whose normalized generators satisfy the following relations with a fixed system of parameters $a = (a(m-k), \ldots, a(m-1))$, w, and z, according to Blackburn [6] and Miech [31, p. 332]:

- parametrized *commutator relations* with parameters $0 \leq a(m-\ell) \leq p-1$ for $1 \leq \ell \leq k$, in particular, with non-vanishing parameter $a(m-k) > 0$,

$$[s_j, y] = \begin{cases} \prod_{\ell=1}^{k-j+2} s_{m-\ell}^{a(m-j+2-\ell)} \in [\gamma_j G, \chi_2 G] = \gamma_{m-k+j-2} G, \text{ for } 2 \leq j \leq k+1, \\ 1, \text{ for } j \geq k+2, \end{cases} \tag{2.10}$$

- parametrized *relations for* pth *powers* of the generators x, y and of higher commutators s_j, with parameters $0 \leq w, z \leq p-1$, and $v := a(m-k)$ if $k = p-2$, but $v := 0$ if $k \leq p-3$,

$$\begin{aligned} (xy^j)^p &= s_{m-1}^{w+zj+vj^2} \in \zeta_1 G, \quad \text{for } 0 \leq j \leq p-1 \\ y^p \prod_{\ell=2}^{p} s_\ell^{\binom{p}{\ell}} &= s_{m-1}^z \in \zeta_1 G, \\ s_{j+1}^p \prod_{\ell=2}^{p} s_{j+\ell}^{\binom{p}{\ell}} &= 1, \quad \text{for } 1 \leq j \leq m-2, \end{aligned} \tag{2.11}$$

- parametrized *nilpotency relations* with parameter $m \geq 3$ (the index of nilpotency),

$$s_j \neq 1, \text{ for } j \leq m-1, \quad \text{but } s_j = 1, \text{ for } j \geq m, \tag{2.12}$$

  expressing the polycyclic structure $\gamma_j G = \langle s_j, \gamma_{j+1} G\rangle$, $\gamma_j G/\gamma_{j+1} G \simeq C_p$, for $j \geq 2$ and $j \leq m-1$, with *cyclic factors* (CF) of the lower central series, which coincides here with the reverse upper central series, $\zeta_j G = \gamma_{m-j} G$, for $0 \leq j \leq m-1$,
- and trivial *metabelian relations* within G',

$$[s_i, s_j] = 1, \text{ for all } i, j \geq 2. \tag{2.13}$$

In the case of an index of nilpotency $m \geq 4$, the commutator relation for s_2 in Formula (2.10) explicitly describes the properties of the two-step centralizer $\chi_2 G = \langle y, \gamma_2 G\rangle$, which are implicitly postulated by Formula (2.8):

$$[\gamma_2 G, \chi_2 G] = \gamma_{m-k} G = \zeta_k G, \tag{2.14}$$

where $k = k(G)$ denotes the *defect of commutativity* of G, which displays the following variety of possible values: $k = 0$ for $3 \leq m \leq 4$, $0 \leq k \leq m-4$ for $m \geq 5$, and $0 \leq k \leq \min(m-4, p-2)$ for $m \geq p+1$, according to Miech [31, p. 331].

The maximal normal subgroups M_i of G contain the commutator subgroup $\gamma_2 G$ of G as a normal subgroup of index p and thus are of the shape $M_i = \langle g_i, \gamma_2 G \rangle$. We define a standard ordering by $g_1 = y$ and $g_i = xy^{i-2}$ for $2 \leq i \leq p+1$. In particular, $M_1 = \langle y, \gamma_2 G \rangle = \chi_2 G$ is the two-step centralizer. In summary:

$$M_1 = \langle y, \gamma_2 G \rangle = \chi_2 G, \quad \text{and} \quad M_i = \langle xy^{i-2}, \gamma_2 G \rangle \quad \text{for } 2 \leq i \leq p+1. \tag{2.15}$$

2.3. **Polarization of the transfer target type.** The transfer target type $\tau(G)$ of a non-abelian p-group G whose second derived quotient G/G'' is of maximal class has a particularly simple form with a single *polarized* component and p fixed components of type $[p,p]$. The polarization is due to the *two-step centralizer* $\chi_2\mathfrak{G}$ of the metabelianization $\mathfrak{G} := G/G''$ of G [25, § 3.1.1, p. 412].

Definition 2.3. Let p be a prime number and $e \geq p-1$ be an integer. By the *nearly homocyclic* abelian p-group of order p^e we understand the abelian group

$$A(p,e) := \left(\overbrace{p^{q+1}, \ldots, p^{q+1}}^{r \text{ times}}, \overbrace{p^q, \ldots, p^q}^{p-1-r \text{ times}} \right), \tag{2.16}$$

with r invariants p^{q+1} and $p-1-r$ invariants p^q, where $q \geq 1$ and $0 \leq r < p-1$ denote the quotient and remainder of the Euclidean division $e = q \cdot (p-1) + r$ of e by $p-1$. Additionally, let

$$A(p,r) := \left(\overbrace{p, \ldots, p}^{r \text{ times}} \right), \tag{2.17}$$

be the *elementary* abelian p-group of rank r for $1 \leq r < p-1$, and $A(p,0) := 1$ the trivial group.

The abelian group $A(p,e)$ is homocyclic if and only if e is a multiple of $p-1$ or $1 \leq e < p-1$.

Theorem 2.2. *Let p be a prime number and G be a p-group whose second derived quotient $\mathfrak{G} := G/G''$ is of coclass $\mathrm{cc}(\mathfrak{G}) = 1$, nilpotency class $c := \mathrm{cl}(\mathfrak{G}) \geq 2$, and defect $k := k(\mathfrak{G})$, but is neither isomorphic to the extra special p-group $G_0^3(0,1)$ of order p^3 and exponent p^2 nor to the Sylow p-subgroup $G_0^{p+1}(1,0) \simeq \mathrm{Syl}_p \mathrm{A}_{p^2}$ of the alternating group of degree p^2, with odd $p \geq 3$. Then the transfer target type of G is given by*

$$\tau(G) = \left(A(p, c-k), \overbrace{[p,p], \ldots, [p,p]}^{p \text{ times}} \right). \tag{2.18}$$

Proof. According to Theorem 6.2, G and $\mathfrak{G}$ share a common restricted Artin pattern, $\mathrm{AP}(G) = \mathrm{AP}(\mathfrak{G})$. In particular they share the same transfer target type $\tau(G) = \tau(\mathfrak{G})$. Thus we have reduced the claim to the metabelian p-group $\mathfrak{G}$ of maximal class, as discussed in § 2.2.

We use the standard ordering of the maximal subgroups $M_i \trianglelefteq \mathfrak{G}$ in Formula (2.15) and recall from [23, Cor. 3.1, p. 476] that the commutator subgroups M_i' are given by

$$M_1' = \gamma_{m-k}\mathfrak{G}, \quad \text{and} \quad M_i' = \gamma_3\mathfrak{G} \text{ for } 2 \leq i \leq p+1. \tag{2.19}$$

We start by showing that $M_i/M_i' \simeq [p,p]$ for $2 \leq i \leq p+1$. To this end, we observe that $M_i = \langle xy^{i-2}, \gamma_2\mathfrak{G} \rangle$ and $\gamma_2\mathfrak{G} = \langle s_2, \gamma_3\mathfrak{G} \rangle$, and thus $M_i/M_i' = \langle xy^{i-2}, s_2, \gamma_3\mathfrak{G} \rangle / \gamma_3\mathfrak{G}$ where x, y are

the normalized generators in Formula (2.9) and $s_2 = [y, x]$ is the main commutator of $\mathfrak{G}$. We have to determine the order of the elements xy^{i-2}, with $2 \le i \le p+1$, and s_2 modulo $\gamma_3\mathfrak{G}$. The third equation in Formula (2.11) yields $s_2^p \cdot \prod_{\ell=2}^{p} s_{1+\ell}^{\binom{p}{\ell}} = 1$ for $j = 1$, and thus $s_2^p \in \gamma_3\mathfrak{G}$, i.e., $\mathrm{ord}(s_2) = p$. The first equation in the same formula gives $(xy^j)^p = s_{m-1}^{w+zj+vj^2}$, for $0 \le j \le p-1$, which is an element of $\gamma_3\mathfrak{G}$ when $m - 1 \ge 3$, that is $m \ge 4$. For $m = 3$, however, we certainly have $k = 0$ and thus $v = 0$, but we must distinguish between the two extra special groups $G_0^3(0, 0)$, where $z = w = 0$ and $(xy^j)^p = 1$, and $G_0^3(0, 1)$, where $z = 0$, $w = 1$ and $(xy^j)^p = s_2$. In summary, we have $\mathrm{ord}(xy^{i-2}) = p^2$ and $M_i/M_i' \simeq [p^2]$ for $G_0^3(0, 1)$, but $\mathrm{ord}(xy^{i-2}) = p$ otherwise, which proves our claim.

We continue by showing that $M_1/M_1' \simeq A(p, c-k)$, where $c = m-1$. In this case, we have $M_1 = \langle y, \gamma_2\mathfrak{G}\rangle$, $\gamma_2\mathfrak{G} = \langle s_2, \ldots, s_{m-k-1}, \gamma_{m-k}\mathfrak{G}\rangle$, and thus $M_1/M_1' = \langle y, s_2, \ldots, s_{m-k-1}, \gamma_{m-k}\mathfrak{G}\rangle/\gamma_{m-k}\mathfrak{G}$. We put $N := m - k$ and use the third equation in Formula (2.11), that is,

$$s_{j+1}^p \cdot s_{j+2}^{\binom{p}{2}} \cdots s_{j+p-1}^{\binom{p}{p-1}} \cdot s_{j+p} = 1, \quad \text{for } 1 \le j \le m-2.$$

These power relations enable us to compute the order of the elements y and s_j, with $2 \le j \le N-1$, modulo $\gamma_N\mathfrak{G}$ by a finite nested double induction with respect to the quotient $q \ge 0$ and the remainder $0 \le r < p-1$ of the Euclidean division $N - 1 = q \cdot (p-1) + r$. Thereby we always have to observe that $s_j \in \gamma_N\mathfrak{G}$, i.e. $\mathrm{ord}(s_j) = 1$, for $j \ge N$, and that the binomial coefficients $\binom{p}{j}$ are multiples of p for $1 \le j \le p-1$.

The first inner induction on the remainders successively yields
$s_{N-1}^p = 1$, $s_{N-2}^p s_{N-1}^{\binom{p}{2}} = 1$, and so on until $s_{N-p+1}^p s_{N-p+2}^{\binom{p}{2}} \cdots s_{N-1}^{\binom{p}{p-1}} = 1$,
that is, we can determine the orders $\mathrm{ord}(s_j) = p$, for $N - (p-1) \le j \le N-1$, recursively.

The second inner induction on the remainders successively yields
$s_{N-p}^p s_{N-1} = 1$, $s_{N-p-1}^p s_{N-p}^{\binom{p}{2}} s_{N-2} = 1$, and so on until $s_{N-2p+2}^p s_{N-2p+3}^{\binom{p}{2}} \cdots s_{N-p}^{\binom{p}{p-1}} s_{N-p+1} = 1$,
that is, we can determine the orders $\mathrm{ord}(s_j) = p^2$, for $N - 2(p-1) \le j \le N - (p-1) - 1$, recursively, but also expressions for s_j with $N - (p-1) \le j \le N-1$ in terms of s_ℓ with $\ell < j$.

In the same manner, we continue to obtain the orders
$\mathrm{ord}(s_j) = p^3$, for $N - 3(p-1) \le j \le N - 2(p-1) - 1$, and so on until

$$\mathrm{ord}(s_j) = p^q, \quad \text{for } N - q(p-1) \le j \le N - (q-1)(p-1) - 1, \tag{2.20}$$

by outer induction on the quotients.

The last inner induction on the remainders will in general be shorter than the previous inner inductions and yields

$$\mathrm{ord}(s_j) = p^{q+1}, \quad \text{for } N - q(p-1) - r \le j \le N - q(p-1) - 1, \tag{2.21}$$

but the recursive process terminates when we reach $N - q(p-1) - (r-1) = 2$, that is,
$s_2^p \cdot s_3^{\binom{p}{2}} \cdots s_p^{\binom{p}{p-1}} \cdot s_{p+1} = 1$,
which shows that the higher commutators $s_{p+1}, \ldots, s_{N-1}$ can be expressed in terms of $s_2, \ldots, s_p$.

Finally we employ the second equation in Formula (2.11), that is,

$$y^p \cdot s_2^{\binom{p}{2}} \cdots s_{p-1}^{\binom{p}{p-1}} \cdot s_p = s_{m-1}^z,$$

which formally corresponds to $N - q(p-1) - r = 1$ when we put $s_1 := y$, but now the right side is s_{m-1}^z instead of 1. Here we must be careful and distinguish the exceptional case of the Sylow p-subgroup $G_0^{p+1}(1, 0) \simeq \mathrm{Syl}_p\mathrm{A}_{p^2}$ of the alternating group of degree p^2, if $p \ge 3$ is an odd prime. In this special case, we have $m = p + 1$, $z = 1$, and thus $s_{m-1}^z = s_p$, which has the unique effect that s_p cancels and cannot be expressed in terms of $y, s_2, \ldots, s_{p-1}$. All elements $y, s_2, \ldots, s_{p-1}, s_p$ are of order p and generate an elementary abelian p-group of rank p, bigger than the rank $p-1$ of all nearly homocyclic abelian p-groups.

In all other cases, the higher commutators $s_p, s_{p+1}, \ldots, s_{N-1}$ can be expressed in terms of $y, s_2, \ldots, s_{p-1}$, and the Formulas (2.20) and (2.21) show that M_1/M_1' is isomorphic to the nearly homocyclic abelian p-group $A(p, c-k)$ of logarithmic order $N-1 = m-k-1 = c-k$. □

Note that Theorem 2.2 is more general than Blackburn's Theorem [6, Thm. 3.4, p. 68], since we have used Theorem 6.2 and therefore the p-group G need not be of maximal class.

3. Monotony of Artin patterns on descendant trees

Definition 3.1. Let p be a prime and G, H and R be finite p-groups.

(1) The *lower central series* (LCS) of G is defined recursively by

$$\gamma_1 G := G, \quad \text{and} \quad \gamma_n G := [\gamma_{n-1} G, G] \text{ for } n \geq 2. \tag{3.1}$$

(2) We call G an *immediate descendant* (or *child*) of H, and H *the parent* of G, if $H \simeq G/\gamma_c G$ is isomorphic to the image of the natural projection $\pi : G \to G/\gamma_c G$ of G onto the quotient by the last non-trivial term $\gamma_c G > 1$ of the LCS of G, where $c := \mathrm{cl}(G)$ denotes the *nilpotency class* of G. In this case, we consider the projection π as a directed edge $G \to H$ from G to $H \simeq \pi G$, and we speak about the *parent operator* π:

$$\pi : G \to \pi G = G/\gamma_c G \simeq H \quad \text{with} \quad c = \mathrm{cl}(G). \tag{3.2}$$

(3) We call G a *descendant* of H, and H an *ancestor* of G, if there exists a finite *path* of directed edges

$$(Q_j \to Q_{j+1})_{0 \leq j < \ell} \quad \text{such that} \quad G = Q_0 \text{ and } H = Q_\ell, \tag{3.3}$$

where $\ell \geq 0$ denotes the *path length*.

(4) The *descendant tree* of R, denoted by $\mathcal{T}(R)$, is the rooted directed tree with root R having the isomorphism classes of all descendants of R as its *vertices* and all (child, parent)-pairs (G, H) among the descendants G, H of R as its *directed edges* $G \to \pi G \simeq H$. By means of formal iterations π^j of the parent operator π, each vertex of the descendant tree $\mathcal{T}(R)$ can be connected with the root R by a finite path of edges:

$$\mathcal{T}(R) = \left\{ G \mid G = \pi^0 G \to \pi^1 G \to \pi^2 G \to \ldots \to \pi^\ell G = R \quad \text{for some} \quad \ell \geq 0 \right\}. \tag{3.4}$$

The restricted Artin pattern enjoys the following monotony property on a descendant tree.

Theorem 3.1. *Let $\mathcal{T}(R)$ be the descendant tree with root $R > 1$, a finite non-trivial p-group, and let $G \to \pi G$ be a directed edge of the tree. Then the restricted Artin pattern $\mathrm{AP} = (\tau, \varkappa)$ satisfies the following monotonicity relations*

$$\begin{aligned} \tau(G) &\geq \tau(\pi G), \\ \varkappa(G) &\leq \varkappa(\pi G), \end{aligned} \tag{3.5}$$

that is, the TTT τ is an isotonic mapping and the TKT $\varkappa$ is an antitonic mapping with respect to the partial order $G > \pi G$ induced by the directed edges $G \to \pi G$.

Proof. An abelian p-group A has the trivial group $\pi A = A/\gamma_c A \simeq 1$ as its parent, because the last non-trivial lower central $\gamma_c A = \gamma_1 A = A$ of A coincides with A. Since the tree root $R > 1$ is supposed to be non-trivial, none of its proper descendants $G > R$ can be abelian. G must be of class $c = \mathrm{cl}(G) \geq 2$. Consequently, if $(G, \pi G)$ is a (child, parent)-pair of the tree $\mathcal{T}(R)$, then the last non-trivial lower central of the child G is contained in the commutator subgroup of G, i.e., $\ker(\pi) = \gamma_c G \leq \gamma_2 G = G'$, as pointed out in [30, Thm. 5.3, p. 85].

The restricted Artin pattern of a finite p-group G is the pair $\mathrm{AP}(G) = (\tau(G), \varkappa(G))$ consisting of the transfer target type $\tau(G) = (U/U')_{G' < U < G}$ and the transfer kernel type $\varkappa(G) =$

$(\ker T_{G,U})_{G'<U<G}$, where $T_{G,U} : G \to U/U'$ denotes the Artin transfer from G to the abelianization U/U' of an intermediate group $G' < U < G$. For each of these intermediate groups, we have $\ker(\pi) \leq G' < U$, which admits several statements about the epimorphism $\pi : G \to \pi G$.

According to [30, Prop.5.1, p. 82], the mapping π is a bijection between the following systems of subgroups, $\mathcal{U} := \{U \mid G' < U < G\}$ and $\mathcal{V} := \{V \mid (\pi G)' = \pi(G') < V = \pi U < \pi G\}$, which is a necessary condition for the uniform comparability of the Artin patterns $\mathrm{AP}(G)$ and $\mathrm{AP}(\pi G)$ of (child, parent)-pairs $(G, \pi G)$ on the tree $\mathcal{T}(R)$.

The TTT τ is an isotonic mapping on the tree $\mathcal{T}(R)$, since $\pi U/\pi U'$ is an epimorphic image of U/U', according to [30, Thm. 5.1, p. 78], and this property was used to define a partial order $\pi U/\pi U' \leq U/U'$ on the components of the TTT, in [30, Dfn. 5.1, p. 80]. Combining all components of the TTT, we obtain $\tau(\pi G) \leq \tau(G)$, by [30, Dfn. 5.4, pp. 83–84]. The (non-strict) inequality has the same direction as $\pi G < G$.

The TKT $\varkappa$ is an antitonic mapping on the tree $\mathcal{T}(R)$, since $\pi(\ker(T_{G,U})) \leq \ker(T_{\pi G,\pi U})$, according to [30, Thm. 5.2, p. 80], and this property was used to define a partial order $\ker(T_{G,U}) \leq \ker(T_{\pi G,\pi U})$ on the components of the TKT, in [30, Dfn. 5.2, p. 82]. Combining all components of the TKT, we obtain $\varkappa(G) \leq \varkappa(\pi G)$, by [30, Dfn. 5.4, pp. 83–84]. The (non-strict) inequality has the opposite direction as $G > \pi G$. □

Remark 3.1. Due to the transitivity of all partial orders involved, Theorem 3.1 remains true, when the (child, parent)-pair $(G, \pi(G))$ is replaced by any (descendant, ancestor)-pair (G, H).

For the proof of our principal results, we require some additional inheritance properties of group theoretic invariants with respect to the (child, parent)-relation of a descendant tree. The inheritance is directed from the child to its parent.

Definition 3.2. A finite or infinite pro-p group G, with an odd prime $p \geq 3$, is called a *σ-group*, if it possesses a *generator inverting* (GI-)automorphism $\sigma \in \mathrm{Aut}(G)$ which acts as the inversion mapping on the derived quotient G/G',

$$\sigma(g)G' = g^{-1}G' \quad \text{for all } g \in G. \tag{3.6}$$

Proposition 3.1. *Let $\pi : G \to \pi G$ be an epimorphism of groups and let $p \geq 2$ be a prime number.*

1. *Let G be a pro-p group, with an odd prime $p \geq 3$.*
   If G is a σ-group, then πG is also a σ-group.
2. *Let G be a finite p-group and let $\pi : G \to \pi G$ be the parent operator $\pi G = G/\gamma_c G$ with $c = \mathrm{cl}(G)$.*
   - *The nilpotency class increases exactly by 1 from the parent πG to the child G.*
   - *If s denotes the step size from parent πG to child G, then the coclass of G is given by $\mathrm{cc}(G) = \mathrm{cc}(\pi G) + (s-1)$, in particular, $\mathrm{cc}(G) = \mathrm{cc}(\pi G)$ for $s = 1$.*
   - *The derived length of πG is not bigger than the derived length of G.*

$$\mathrm{cl}(G) = \mathrm{cl}(\pi G) + 1, \quad \mathrm{cc}(G) = \mathrm{cc}(\pi G) + (s-1), \qquad \mathrm{dl}(G) \geq \mathrm{dl}(\pi G). \tag{3.7}$$

3. *If G is a finite metabelian p-group of maximal class and π is the parent operator, then the defect of the parent πG is not bigger than the defect of the child G, that is, $k(\pi G) \leq k(G)$, more precisely*

$$k(\pi G) = \begin{cases} k(G) & \text{if } k(G) = 0, \\ k(G) - 1 & \text{if } k(G) \geq 1, \end{cases} \tag{3.8}$$

which can also be expressed in the form $k(\pi G) = \max(0, k(G) - 1)$.

Proof. Let $d := \mathrm{dl}(G)$ be the derived length of G and

$$G = G^{(0)} > G^{(1)} > \ldots > G^{(d-1)} > G^{(d)} = 1$$

be the derived series of G, defined recursively by $G^{(0)} := G$ and $G^{(j)} := [G^{(j-1)}, G^{(j-1)}]$ for $j \geq 1$. For any homomorphism $\pi : G \to H$ of groups, the proof of [30, Cor. 7.2, Appendix, p. 100] has shown that $(\pi G)^{(j)} = \pi\left(G^{(j)}\right)$ for $j \geq 0$, and similarly for the lower central series $\gamma_j(\pi G) = \pi\left(\gamma_j G\right)$ for $j \geq 1$.

(1) The inheritance of a GI-automorphism has been proved in [30, Thm. 7.2, App., p. 102].

(2) The lower central series of πG is given by
$\pi G > \gamma_2\pi G = \pi\gamma_2 G = \gamma_2 G/\gamma_c G > \ldots > \gamma_{c-1}\pi G = \pi\gamma_{c-1}G = \gamma_{c-1}G/\gamma_c G > \gamma_c\pi G \simeq 1$,
whence $\mathrm{cl}(\pi G) = c - 1$.

Let n be the logarithmic order of G, that is $|G| = p^n$. If $|\gamma_c G| = p^s$, then we have $|\pi G| = (G : \gamma_c G) = p^{n-s}$ and πG is of logarithmic order $n - s$. Consequently, the coclass of πG is given by $\mathrm{cc}(\pi G) = (n - s) - (c - 1) = (n - c) - (s - 1) = \mathrm{cc}(G) - (s - 1)$.

The image of the last non-trivial term $G^{(d-1)}$ of the derived series of G under π is given by $\pi G^{(d-1)} = G^{(d-1)} \ker\pi/\gamma_c G$, where $\ker\pi = \gamma_c G$. If we denote by $d' := \mathrm{dl}(\pi G)$ the derived length of πG, then the derived series of πG is given by

$$\pi G = \pi G^{(0)} > \pi G^{(1)} > \ldots > \pi G^{(d'-1)} > \pi G^{(d')} = 1,$$

where $d' < d$ if $G^{(d-1)} \leq \gamma_c G$, and $d' = d$ otherwise.

(3) If the order of G is given by $|G| = p^m$, then the two-step centralizer of G satisfies the condition $[\chi_2 G, \gamma_2 G] = \gamma_{m-k} G$, where $k := k(G) \leq m - 4$ denotes the defect of G. Application of the epimorphism π yields

$$[\chi_2\pi G, \gamma_2\pi G] = [\pi\chi_2 G, \pi\gamma_2 G] = \pi[\chi_2 G, \gamma_2 G] = \pi\gamma_{m-k} G = \gamma_{m-k}\pi G,$$

where

$$\gamma_{m-k}\pi G = \begin{cases} 1 & \text{if } k = 0, \\ \gamma_{(m-1)-(k-1)}\pi G & \text{if } k \geq 1. \end{cases}$$

This proves the claim, since the step size for coclass 1 is $s = 1$ and thus $|\pi G| = p^{m-1}$.
Here, we have used that $\pi\chi_2 G = \chi_2\pi G$, since applying the surjection π to the equation

$$\chi_2 G = \{g \in G \mid [g, u] \in \gamma_4(G) \quad \text{for all } u \in \gamma_2(G)\}$$

and observing that $\pi\gamma_2 G = \gamma_2\pi G$ and $\pi\gamma_4 G = \gamma_4\pi G$ yields

$$\pi\chi_2 G = \{\pi g \in \pi G \mid [\pi g, \pi u] \in \gamma_4\pi G \quad \text{for all } \pi u \in \gamma_2\pi G\}. \qquad \square$$

4. Proof of the main results

We are now in the position to prove Theorem 1.1 and Corollary 1.1 simultaneously. The dominant part of the proof is valid for *any* number field K and admits *two possibilities* $\ell_5 K \in \{2, 3\}$ for the length of its 5-class tower. The final specialization to a real quadratic field $K = \mathbb{Q}(\sqrt{d})$ with discriminant $d > 0$ will enforce the fixed length $\ell_5 K = 3$.

Proof. Let K be an *arbitrary* algebraic number field with 5-class group $\mathrm{Cl}_5 K \simeq [5, 5]$ and 5-capitulation type $\varkappa(K) = \left(\ker j_{L_i|K}\right)_{1 \leq i \leq 6} \sim (1, 0^5)$ in its six unramified cyclic quintic extensions $L_1, \ldots, L_6$, and suppose that the 5-class groups of the latter are given by $\tau(K) = \left(\mathrm{Cl}_5 L_i\right)_{1 \leq i \leq 6} \sim \left([25, 5, 5, 5], [5, 5]^5\right)$.

Then the restricted Artin pattern of K is $\mathrm{AP}(K) = (\tau(K), \varkappa(K))$ and, by Theorem 2.1, Formula (2.5), which constitutes the translation from number theory to group theory, $\mathrm{AP}(K)$ coincides with the restricted Artin pattern $\mathrm{AP}(\mathfrak{G}) = (\tau(\mathfrak{G}), \varkappa(\mathfrak{G}))$ of the second 5-class group $\mathfrak{G} = \mathrm{G}_5^2 K$, where $\tau(\mathfrak{G}) = \left(M_i/M_i'\right)_{1 \leq i \leq 6}$ and $\varkappa(\mathfrak{G}) = \left(\ker T_{\mathfrak{G}, M_i}\right)_{1 \leq i \leq 6}$ in terms of the six maximal subgroups $M_1, \ldots, M_6$ of $\mathfrak{G}$ and the Artin transfers $T_{\mathfrak{G}, M_i}$.

According to Formula (1.2), $\mathfrak{G}$ possesses the abelianization $\mathfrak{G}/\mathfrak{G}' \simeq \mathrm{Cl}_5 K \simeq [5, 5]$. Hence, the metabelian 5-group $\mathfrak{G}$ must be a descendant of the abelian 5-group $\langle 5^2, 2\rangle \simeq [5, 5]$, and for identifying $\mathfrak{G}$ by the strategy of *pattern recognition via Artin transfers* we therefore have to start the p-group generation algorithm [18] at the root $R = \langle 5^2, 2\rangle$ and to construct the descendant tree $\mathcal{T}(R)$, always looking for the *assigned search pattern* $\mathrm{AP}(\mathfrak{G})$.

The recursive steps of this algorithm by Newman [33] and O'Brien [35] may be monitored on the metabelian skeleton of the coclass-1 subtree $\mathcal{T}^1(R)$ in [25, Fig. 3.3, p. 425], on the interface between coclass 1 and 2 in [25, Fig. 3.8, p. 448], and on the pruned descendant tree containing the decisive bifurcation in Figure 1.

In each step, a parent $\pi(G)$ gives rise to several children G, and the TTT $\tau(G) \geq \tau(\pi(G))$ is isotonic with respect to the partial order $G > \pi(G)$ induced by the tree edges $G \to \pi(G)$, wheras the TKT $\varkappa(G) \leq \varkappa(\pi(G))$ is antitonic, by the *fundamental monotony result* in Theorem 3.1.

The *pruning* process will run parallel to the recursive tree construction in the following way. A vertex with a first TTT component $\tau_1 > [25,5,5,5]$ or with a higher TTT component $\tau_i > [5,5]$, for some $2 \leq i \leq 6$, must be discarded together with all its descendants, since their TTT τ will contain an inadequate component. Similarly, a vertex having any *forbidden* TKT $\varkappa$ different from the only two *admissible* TKTs, $\varkappa \sim (1,0^5)$ and the biggest possible $\varkappa = (0^6)$, must be cancelled together with the entire set of its descendants, since their TKT $\varkappa$ will also be forbidden.

An important *additional filter* must be used to eliminate the numerous metabelian 5-groups of maximal class with positive defect of commutativity, $k \geq 1$, although they share the common admissible TKT $\varkappa = (0^6)$, by [24, Thm. 2.5.(3), p. 479]. However, all descendants of a parent with $k \geq 1$ will also have $k \geq 1$, by item (3) of Proposition 3.1, Formula (3.8), and a metabelian group of maximal class with $k \geq 1$ reveals a *total stabilization* of all components of the restricted Artin pattern, according to Theorem [30, Thm. 6.2.(2), p. 92]. With the aid of Theorem 6.2, Formula (6.4), we conclude that non-metabelian 5-groups G of any coclass, whose metabelianization G/G'' is of maximal class with $k \geq 1$, will also have the TKT $\varkappa = (0^6)$ (and not the desired TKT $\varkappa \sim (1,0^5)$) and, due to $k \geq 1$, the inadequate first TTT component which was determined in Theorem 2.2. (A metabelian p-group of maximal class with order p^m has defect $k \geq 1$ if and only if $|\tau_1| < m-1$.)

After these preliminary considerations, we execute the actual steps for finding *successive approximations* in form of *class-c quotients* $G/\gamma_{c+1}G$ for the 5-class tower group $G := \mathrm{G}_5^\infty K$ of the field K. The steps are summarized in Table 1.

- The abelian root $R = \langle 5^2, 2\rangle \simeq C_5 \times C_5$ is the class-1 quotient $G/\gamma_2 G$ of G, as shown by Formula (1.2). It has $\tau = \big([5]^6\big)$, $\varkappa = (0^6)$, and gives rise to the extra special 5-groups as its 2 children, the capable $\langle 5^3, 3\rangle \simeq G_0^3(0,0)$ with $\tau = \big([5,5]^6\big)$, and admissible $\varkappa = (0^6)$, and the terminal $\langle 5^3, 4\rangle \simeq G_0^3(0,1)$ with $\tau \sim \big([5,5],[25]^5\big)$, and forbidden $\varkappa = (1^6)$.
  (For a real quadratic field K, the group $\langle 5^3, 4\rangle$ cannot occur as $\mathrm{G}_5^2 K$ for the additional reason that it is not a σ-group.)
- The compulsory next parent $\langle 5^3, 3\rangle \simeq G_0^3(0,0)$ is the class-2 quotient $G/\gamma_3 G$ of G and reveals a *bifurcation* [26, § 8, p. 168] with 4 metabelian children of step size 1 [25, Fig. 3.3, p. 425], and 12 metabelian children of step size 2 [25, Fig. 3.8, p. 448].

  Among the former, we have a capable vertex $\langle 5^4, 7\rangle$ with admissible $\varkappa = (0^6)$, a single leaf $\langle 5^4, 8\rangle$ with the desired $\varkappa \sim (1,0^5)$ but too small $\tau \sim \big([5,5,5],[5,5]^5\big)$, and two leaves $\langle 5^4, 9|10\rangle$ with forbidden $\varkappa \sim (2,0^5)$.

  Among the latter, which are of coclass 2, the 11 groups $\langle 5^5, 4 \ldots 14\rangle$ have forbidden TKTs, and only the group $\langle 5^5, 3\rangle$ has admissible $\varkappa = (0^6)$. However, the last five components of the TTT $\tau = \big([5,5,5]^6\big)$ disqualify the group as a possible class-3 quotient of G.
- For the class-3 quotient $G/\gamma_4 G$ of G we have to take $\langle 5^4, 7\rangle \simeq G_0^4(0,0)$ which is the next parent. It has 9 metabelian children, all of step size 1. The capable vertex $\langle 5^5, 30\rangle$ has admissible $\varkappa = (0^6)$. The leaf $\langle 5^5, 31\rangle$ has the desired $\varkappa \sim (1,0^5)$ but too small $\tau \sim \big([5,5,5,5],[5,5]^5\big)$. The leaf $\langle 5^5, 32\rangle$ has forbidden $\varkappa \sim (2,0^5)$, and the 6 groups $\langle 5^5, 33 \ldots 38\rangle$ have positive defect of commutativity $k = 1$.
  (For a real quadratic field K, the groups $\langle 5^5, 31 \ldots 38\rangle$ cannot occur as $\mathrm{G}_5^2 K$ for the additional reason that they are not σ-groups.)
- Now we arrive at the *crucial bifurcation* [26, § 8, p. 168] of the mandatory next parent $\langle 5^5, 30\rangle \simeq G_0^5(0,0)$ which is the class-4 quotient $G/\gamma_5 G$ of G. It has 21 children of step size 1 (13 of them metabelian) and 115 non-metabelian children of step size 2.

Among the former, we have a capable vertex $\langle 5^6, 630\rangle$ with admissible $\varkappa = (0^6)$, 4 vertices $\langle 5^6, 631 \dots 634\rangle$ with forbidden $\varkappa \sim (2, 0^5)$, a single group $\langle 5^6, 635\rangle$ with both, the desired $\varkappa \sim (1, 0^5)$ and the desired $\tau \sim \big([25, 5, 5, 5], [5, 5]^5\big)$, and 7 groups $\langle 5^6, 636 \dots 642\rangle$ with $k = 1$.

Among the latter, which are of coclass 2, we have 3 capable vertices $\langle 5^7, 360|372|384\rangle$ with admissible $\varkappa = (0^6)$, 5 leaves $\langle 5^7, 361|373|374|385|386\rangle$ with both, the desired $\varkappa \sim (1, 0^5)$ and the desired $\tau \sim \big([25, 5, 5, 5], [5, 5]^5\big)$, 32 vertices $\langle 5^7, n\rangle$, where n is listed in Table 1, with forbidden $\varkappa \sim (2, 0^5)$, and 75 vertices $\langle 5^7, 392 \dots 466\rangle$ with $k \geq 1$.

- To make sure that $\langle 5^6, 635\rangle \simeq G_0^5(0, 1)$ is the unique possibility for the metabelianization G/G'' of G, we have to check the next parent $\langle 5^6, 630\rangle \simeq G_0^5(0, 0)$. It has 12 children of step size 1 (9 of them metabelian), but either their $\tau \sim \big([25, 25, 5, 5], [5, 5]^5\big)$ is too big or they have $k = 1$. Thus, the break-off condition on the coclass-1 tree is reached.
- It remains to prove that $\langle 5^7, 361|373|374|385|386\rangle$ are the only 5 non-metabelian possibilities for G. For this purpose, we must check the roots $\langle 5^7, 360|372|384\rangle$ of 3 non-metabelian coclass-2 trees, which possess 14, resp. 20, resp. 20, children of step size 1, but either their $\tau \sim \big([25, 25, 5, 5], [5, 5]^5\big)$ is too big or they have $k \geq 1$.

The proof for a real quadratic field K will be finished in Theorem 6.3. □

TABLE 1. Descendants of class-c quotients $G/\gamma_{c+1}G$ of the 5-class tower group G

$G/\gamma_{c+1}G$				step	$\varkappa = (0^6)$ $k = 0$		$\varkappa \sim (1, 0^5)$ $k = 0$		$\varkappa \sim (2, 0^5)$ $k = 0$		$\varkappa = (0^6)$ $k = 1$	
c	ord	id	# of descendants	size	#	id	#	id	#	id	#	id
1	5^2	2	2/1	1	1	3						
2	5^3	3	4/1; 12/6	1	1	7	1	8	2	9|10		
				2	1	3						
3	5^4	7	9/2	1	1	30	1	31	1	32	6	33 … 38
4	5^5	30	21/2; 115/13	1	1	630	1	635	4	631 … 634	15	636 … 650
				2	3	360	5	361	32	352 … 359	75	392 … 466
						372		373|374		362 … 371		
						384		385|386		375 … 383		
										387 … 391		
	5^6	630	12/2	1	1	1287	1	1288	1	1286	9	1289 … 1297
	5^7	360	14/2	1	1	2	1	4	2	1|3	10	5 … 14
	5^7	372	20/2	1	1	2	1	5	3	1|3|4	15	6 … 20
	5^7	384	20/2	1	1	2	1	5	3	1|3|4	15	6 … 20

Remark 4.1. It is very important to point out that, for the sake of homogeneity, we have conducted both, the search for the metabelianization $\mathfrak{G} = G/G''$ of the 5-class tower group G, and the search for G itself, within the frame of the p-group generation algorithm. In two previous papers, we have developed the tools for finding the second 5-class group $\mathfrak{G}$ much more directly: The metabelian skeleton of the coclass tree $\mathcal{T}^1(\langle 25, 2\rangle)$ in [25, Fig. 3.3, p. 425] is embedded within a sort of coordinate system with the TKT $\varkappa$ as its horizontal axis and the polarized first TTT component τ_1 as its vertical axis. According to [24, Thm. 2.5.(3), p. 479], the TKT $\varkappa(\mathfrak{G}) \sim (1, 0^5)$ uniquely determines the parameters $w = 1$, $z = 0$, and $k = 0$ of the metabelian 5-group $\mathfrak{G} \simeq G_a^m(z, w) = G_0^m(0, 1)$, since $k = 0$ if and only if a is the empty family. As second input data we do not even need the exact TTT $\tau(\mathfrak{G}) = \big([25, 5, 5, 5], [5, 5]^5\big)$. The order of the polarized first TTT component $|\tau_1| = 25 \cdot 5 \cdot 5 \cdot 5 = 5^5$ suffices, since according to [23, Thm. 3.1.(3), p. 475], the relation $|\tau_1| = |M_1/M_1'| = 5^{m-k-1} = 5^{m-1}$ uniquely determines the index of nilpotency as $m = 6$. Therefore, we have $\mathfrak{G} \simeq G_0^6(0, 1)$, and this is exactly the group $\langle 5^6, 635\rangle$.

5. Relation rank and generator rank

Theorem 5.1. *(Shafarevich) Let p be a prime number and denote by ζ a primitive pth root of unity. Let K be a number field with signature (r_1, r_2) and torsionfree Dirichlet unit rank $r = r_1 + r_2 - 1$, and let S be a finite set of non-archimedean or real archimedean places of K. Assume that no place in S divides p.*

Then the relation rank $d_2G_S := \dim_{\mathbb{F}_p} H^2(G_S, \mathbb{F}_p)$ of the Galois group $G_S := \mathrm{Gal}(K_S|K)$ of the maximal pro-p extension K_S of K which is unramified outside of S is bounded from above by

$$d_2G_S \leq \begin{cases} d_1G_S + r & \text{if } S \neq \emptyset \text{ or } \zeta \notin K, \\ d_1G_S + r + 1 & \text{if } S = \emptyset \text{ and } \zeta \in K, \end{cases} \tag{5.1}$$

where $d_1G_S := \dim_{\mathbb{F}_p} H^1(G_S, \mathbb{F}_p)$ denotes the generator rank of G_S.

Proof. The original statement in [36, Thm. 6, (18′)] contained a serious misprint which was corrected in [29, Thm. 5.5, p. 28]. □

6. Covers and Shafarevich covers

We begin with a purely group theoretic concept, which aids in getting an overview of all extensions of a particular kind of an assigned p-group.

Definition 6.1. Let p be a prime and $\mathfrak{G}$ be a finite *metabelian* p-group. By the *cover* of $\mathfrak{G}$ we understand the set of all (isomorphism classes of) finite p-groups whose second derived quotient is isomorphic to $\mathfrak{G}$:

$$\mathrm{cov}(\mathfrak{G}) := \{G \mid \mathrm{ord}(G) < \infty,\ G/G'' \simeq \mathfrak{G}\}. \tag{6.1}$$

By eliminating the finiteness condition, we obtain the *complete cover* of $\mathfrak{G}$,

$$\mathrm{cov}_c(\mathfrak{G}) := \{G \mid G/G'' \simeq \mathfrak{G}\}. \tag{6.2}$$

Remark 6.1. The unique metabelian element of $\mathrm{cov}(\mathfrak{G})$ is the isomorphism class of $\mathfrak{G}$ itself.

Theorem 6.1. *The cover of the metabelian 5-group $\langle 15625, 635\rangle$ consists of 6 elements,*

$$\mathrm{cov}(\langle 15625, 635\rangle) = \Big\{\langle 15625, 635\rangle\Big\} \bigcup \Big\{\langle 78125, n\rangle \mid n \in \{361, 373, 374, 385, 386\}\Big\}. \tag{6.3}$$

Proof. This is a consequence of the presentations in Formulas (7.1) and (7.2) together with § 4. □

Remark 6.2. We shall not make essential use of the complete cover in this article. The only reason for its introduction is the possibility to express certain assertions in a more general way.

Theorem 6.2. *Let $\mathfrak{G}$ be a finite metabelian p-group. Then all elements of the complete cover of $\mathfrak{G}$ share a common restricted Artin pattern:*

$$\mathrm{AP}(G) = \mathrm{AP}(\mathfrak{G}), \quad \text{for all} \quad G \in \mathrm{cov}_c(\mathfrak{G}). \tag{6.4}$$

Proof. This is the Main Theorem of [30, Thm. 5.4, p. 86]. □

Motivated by the Shafarevich Theorem 5.1, we connect the group theoretic cover concept with arithmetical information, which will be useful for identifying the p-class tower group of an assigned number field.

Definition 6.2. Let p be a prime and K be a number field with p-class rank $\varrho := d_1(\mathrm{Cl}_p(K))$, torsionfree Dirichlet unit rank r, and the second p-class group $\mathfrak{G} := \mathrm{G}_p^2K$. By the *Shafarevich cover*, $\mathrm{cov}(\mathfrak{G}, K)$, of $\mathfrak{G}$ *with respect to* K we understand the subset of $\mathrm{cov}(\mathfrak{G})$ whose elements G satisfy the following condition for their relation rank:

$$\varrho \le d_2 G \le \varrho + r + \vartheta, \quad \text{where} \quad \vartheta := \begin{cases} 1 & \text{if } K \text{ contains the } p\text{th roots of unity,} \\ 0 & \text{otherwise.} \end{cases} \tag{6.5}$$

Theorem 6.3. *The Shafarevich cover of the metabelian* 5*-group* $\langle 15625, 635 \rangle$ *with respect to a real quadratic field* K *of* 5*-class rank* $\varrho = 2$ *consists of* 5 *elements,*

$$\operatorname{cov}(\langle 15625, 635 \rangle, K) = \Big\{ \langle 78125, n \rangle \mid n \in \{361, 373, 374, 385, 386\} \Big\}. \tag{6.6}$$

Proof. A real quadratic field has torsionfree unit rank $r = 0 + 2 - 1 = 1$ and does not contain the primitive fifth roots of unity, i.e., $\vartheta = 0$. The relation rank of a p-group is equal to its p-multiplicator rank [26, (26), p. 178]. The metabelian 5-group $\mathfrak{G} = \langle 15625, 635 \rangle$ has relation rank 4, whereas the relation rank of the non-metabelian 5-groups $G = \langle 78125, n \rangle$ with $n \in \{361, 373, 374, 385, 386\}$ is 3. Consequently, the former satisfies the conflicting condition $d_2\mathfrak{G} = 4 > \varrho + r + \vartheta = 3$, whereas the latter enjoy the required property $2 = \varrho \le d_2 G = 3 \le \varrho + r + \vartheta = 3$ for a 5-class tower group of K. □

Remark 6.3. The arithmetical invariants $\tau(K)$ and $\varkappa(K)$ of the real quadratic fields K in Example 1.1 were computed with the aid of MAGMA, version V2.21-11 [7, 8, 22], under the LINUX operating system on a machine with two XEON 8-core CPUs and 256 GB RAM. The unramified cyclic quintic extensions were constructed by means of the class field routines written by Fieker [12] for MAGMA.

7. Presentation and normal lattice of the 5-tower groups

The diagram in Figure 1 visualizes the tree of 5-groups G with abelianization G/G' of type $[5, 5]$ up to order 5^{12}, *pruned* by restriction to the transfer kernel types a.1, $\varkappa = (0, 0, 0, 0, 0, 0)$, of vertices with defect $k = 0$ on infinite paths, and a.2, $\varkappa \sim (1, 0, 0, 0, 0, 0)$, of leaves.

Groups of coclass $\operatorname{cc}(G) = 1$ are metabelian, and groups of coclass $\operatorname{cc}(G) = 2$, except $\langle 3125, 3 \rangle$, have derived length $\operatorname{dl}(G) = 3$. The big contour square □ denotes the abelian root of the tree, full discs • denote metabelian groups with cyclic centre of order 5, a contour circle ∘ denotes a metabelian group with bicyclic centre of type $(5, 5)$, and smaller contour squares □ denote non-metabelian groups. The symbol 2^* means a batch of two siblings with common parent, where only one vertex is drawn

Up to order 5^7, vertices are labelled by their SmallGroups Library identifiers [4, 5] in angle brackets, where we omit the order, which is given on the left hand scale. Starting with order 5^8, relative identifiers in the form $-\#s; n$, with step size s and counter n, are used as in the output produced by the ANUPQ package [14] of GAP [15] and MAGMA [22]. For additional ease of identification, we give the descendant numbers of some relevant capable vertices in the format $(N_1/C_1; \ldots; N_\nu/C_\nu)$ as described in [26, (34), p.180], where ν denotes the nuclear rank.

The diagram in Figure 2 visualizes the lattice of all normal subgroups of the 5-class tower groups $G_n := \langle 78125, n \rangle$ with $n \in \{361, 373, 374, 385, 386\}$. Polycyclic power commutator presentations of these groups are given by

$$\begin{aligned} G_n = \langle\ & x, y, s_2, s_3, s_4, s_5, u_5 \ \mid \\ & s_2 = [y, x],\ s_3 = [s_2, x],\ s_4 = [s_3, x],\ s_5 = [s_4, x], \\ & x^5 = W,\ y^5 = s_5^4,\ u_5 = [s_3, y] = [s_4, y],\ [s_3, s_2] = u_5^4\ \rangle, \end{aligned} \tag{7.1}$$

where the relator word W for the fifth power of the first generator x is decisive:

$$W = \begin{cases} s_5 & \text{if } n = 361, \\ s_5 u_5 & \text{if } n = 373, \\ s_5^2 u_5 & \text{if } n = 374, \\ s_5 u_5^2 & \text{if } n = 385, \\ s_5^2 u_5^2 & \text{if } n = 386. \end{cases} \tag{7.2}$$

FIGURE 1. Pruned descendant tree of the abelian root $\langle 25, 2 \rangle$



FIGURE 2. Normal lattice of the 5-tower groups $\langle 78125, n\rangle$ ($n \in \{361, 373, 374, 385, 386\}$)



The presentation for the second derived quotient of G_n is obtained by putting $u_5 := 1$. Both resulting relator words $W = s_5$ and $W = s_5^2$ yield the same metabelian group $\langle 15625, 635\rangle$.

8. Acknowledgements

The author gratefully acknowledges that his research is supported by the Austrian Science Fund (FWF): P 26008-N25.

References

[1] E. Artin, *Beweis des allgemeinen Reziprozitätsgesetzes*, Abh. Math. Sem. Univ. Hamburg **5** (1927), 353–363, DOI 10.1007/BF02952531.

[2] E. Artin, *Idealklassen in Oberkörpern und allgemeines Reziprozitätsgesetz*, Abh. Math. Sem. Univ. Hamburg **7** (1929), 46–51, DOI 10.1007/BF02941159.

[3] Y. Berkovich, *Groups of prime power order, Vol. 1*, De Gruyter Expositions in Mathematics, Vol. **46**, Walter de Gruyter, Berlin, New York, 2008.

[4] H. U. Besche, B. Eick and E. A. O'Brien, *A millennium project: constructing small groups*, Int. J. Algebra Comput. **12** (2002), 623-644, DOI 10.1142/s0218196702001115.

[5] H. U. Besche, B. Eick, and E. A. O'Brien, *The SmallGroups Library — a Library of Groups of Small Order*, 2005, an accepted and refereed GAP package, available also in MAGMA.

[6] N. Blackburn, *On a special class of p-groups*, Acta Math. **100** (1958), 45–92.

[7] W. Bosma, J. Cannon, and C. Playoust, *The Magma algebra system. I. The user language*, J. Symbolic Comput. **24** (1997), 235–265.

[8] W. Bosma, J. J. Cannon, C. Fieker, and A. Steels (eds.), *Handbook of Magma functions* (Edition 2.21, Sydney, 2015).

[9] N. Boston and H. Nover, *Computing pro-p Galois groups*, Proceedings of ANTS 2006, Lecture Notes in Computer Science **4076**, 1–10, Springer-Verlag Berlin Heidelberg, 2006.

[10] M. R. Bush, *Computation of Galois groups associated to the* 2*-class towers of some quadratic fields*, J. Number Theory **100** (2003), 313–325.

[11] M. R. Bush and D. C. Mayer, 3*-class field towers of exact length* 3, J. Number Theory **147** (2015), 766–777, DOI 10.1016/j.jnt.2014.08.010.

[12] C. Fieker, *Computing class fields via the Artin map*, Math. Comp. **70** (2001), no. 235, 1293–1303.

[13] Ph. Furtwängler, *Beweis des Hauptidealsatzes für die Klassenkörper algebraischer Zahlkörper*, Abh. Math. Sem. Univ. Hamburg **7** (1929), 14–36.

[14] G. Gamble, W. Nickel and E. A. O'Brien, *ANU p-Quotient — p-Quotient and p-Group Generation Algorithms*, 2006, an accepted GAP package, available also in MAGMA.

[15] The GAP Group, *GAP – Groups, Algorithms, and Programming — a System for Computational Discrete Algebra*, Version 4.7.8, Aachen, Braunschweig, Fort Collins, St. Andrews, 2015, `(\protect\vrule width0pt\protect\href{http://www.gap-system.org}{http://www.gap-system.org})`.

[16] E. S. Golod and I. R. Shafarevich, *On the class field tower* (Russian), Izv. Akad. Nauk SSSR Ser. Mat. **28** (1964), 261–272. (English transl. in Amer. Math. Soc. Transl. (2) **48** (1965), 91–102.)

[17] H. Hasse, *Bericht über neuere Untersuchungen und Probleme aus der Theorie der algebraischen Zahlkörper. Teil II: Reziprozitätsgesetz*, Jber. der DMV **6** (1930), 1–204

[18] D. F. Holt, B. Eick and E. A. O'Brien, *Handbook of computational group theory*, Discrete mathematics and its applications, Chapman and Hall/CRC Press, 2005.

[19] B. Huppert, *Endliche Gruppen I*, Grundlehren der mathematischen Wissenschaften, Vol. **134**, Springer-Verlag, Berlin, Heidelberg, New York, 1979.

[20] H. Kisilevsky, *Number fields with class number congruent to* 4 *mod* 8 *and Hilbert's theorem* 94, J. Number Theory **8** (1976), 271–279.

[21] H. Koch und B. B. Venkov, *Über den p-Klassenkörperturm eines imaginär-quadratischen Zahlkörpers*, Astérisque **24–25** (1975), 57–67.

[22] The MAGMA Group, *MAGMA Computational Algebra System*, Version 2.21-11, Sydney, 2016, `(\protect\vrule width0pt\protect\href{http://magma.maths.usyd.edu.au}{http://magma.maths.usyd.edu.au})`.

[23] D. C. Mayer, *The second p-class group of a number field*, Int. J. Number Theory **8** (2012), no. 2, 471–505, DOI 10.1142/S1793042112500251X.

[24] D. C. Mayer, *Transfers of metabelian p-groups*, Monatsh. Math. **166** (2012), no. 3–4, 467–495, DOI 10.1007/s00605-010-0277-x.

[25] D. C. Mayer, *The distribution of second p-class groups on coclass graphs*, Journal de Théorie des Nombres de Bordeaux **25** (2013), no. 2, 401–456, DOI 10.5802/jtnb.842.

[26] D. C. Mayer, *Periodic bifurcations in descendant trees of finite p-groups*, Adv. Pure Math. **5** (2015), no. 4, 162–195, DOI 10.4236/apm.2015.54020, Special Issue on Group Theory, March 2015.

[27] D. C. Mayer, *Index-p abelianization data of p-class tower groups*, Adv. Pure Math. **5** (2015) no. 5, 286–313, DOI 10.4236/apm.2015.55029, Special Issue on Number Theory and Cryptography, April 2015.

[28] D. C. Mayer, *Periodic sequences of p-class tower groups*, J. Appl. Math. Phys. **3** (2015), no. 7, 746–756, DOI 10.4236/jamp.2015.37090.

[29] D. C. Mayer, *New number fields with known p-class tower*, Tatra Mountains Math. Publ. **64** (2015), 21 – 57, DOI 10.1515/tmmp-2015-0040, Special Issue on Number Theory and Cryptology '15.

[30] D. C. Mayer, *Artin transfer patterns on descendant trees of finite p-groups*, Adv. Pure Math. **6** (2016), no. 2, 66 – 104, DOI 10.4236/apm.2016.62008, Special Issue on Group Theory Research, January 2016.

[31] R. J. Miech, *Metabelian p-groups of maximal class*, Trans. Amer. Math. Soc. **152** (1970), 331–373.

[32] K. Miyake, *Algebraic investigations of Hilbert's Theorem* 94*, the principal ideal theorem and the capitulation problem*, Expo. Math. **7** (1989), 289–346.

[33] M. F. Newman, *Determination of groups of prime-power order*, pp. 73–84, in: Group Theory, Canberra, 1975, Lecture Notes in Math., vol. **573**, Springer, Berlin, 1977.

[34] H. Nover, *Computation of Galois groups associated to the* 2*-class towers of some imaginary quadratic fields with* 2*-class group* $C_2 \times C_2 \times C_2$, J. Number Theory **129** (2009), 231–245.

[35] E. A. O'Brien, *The p-group generation algorithm*, J. Symbolic Comput. **9** (1990), 677–698, DOI 10.1016/S0747-7171(80)80082-X.

[36] I. R. Shafarevich, *Extensions with prescribed ramification points* (Russian), Publ. Math., Inst. Hautes Études Sci. **18** (1964), 71–95. (English transl. by J. W. S. Cassels in Amer. Math. Soc. Transl., II. Ser., **59** (1966), 128–149.)

[37] A. Scholz und O. Taussky, *Die Hauptideale der kubischen Klassenkörper imaginär quadratischer Zahlkörper: ihre rechnerische Bestimmung und ihr Einfluß auf den Klassenkörperturm*, J. Reine Angew. Math. **171** (1934), 19–41.

Naglergasse 53, 8010 Graz, Austria
E-mail address: algebraic.number.theory@algebra.at
URL: http://www.algebra.at